Limitations of the Green's function approach to analyzing spatial radiative feedbacks

Jonah Bloch-Johnson, Maria Rugenstein, Jonathan Gregory CFMIP 2019 - 1.10.19

multiple regression


Limitations of the Green's function approach to analyzing spatial radiative feedbacks


Jonah Bloch-Johnson, Maria Rugenstein, Jonathan Gregory CFMIP 2019 - 1.10.19

Multiple regression approach is a method for estimating spatial feedbacks from internal variability

Multiple regression approach is a method for estimating spatial feedbacks from internal variability


Long-term goal:
observationally-based
forecast of warming that
accounts for the pattern effect





$$c_1 \frac{dT_1}{dt} = N_1 - H + F_{surf,1}$$

$$c_2 \frac{dT_2}{dt} = N_2 + H + F_{surf,2}$$




$$c_1 \frac{dT_1}{dt} = N_1 - H + F_{surf,1}$$

$$c_2 \frac{dT_2}{dt} = N_2 + H + F_{surf,2}$$

$$N_1 = \lambda_{1,1}T_1 + \lambda_{1,2}T_2 + F_{TOA,1} + F_{CO_2,1}$$

$$N_2 = \lambda_{2,1}T_1 + \lambda_{2,2}T_2 + F_{TOA,2} + F_{CO_2,2}$$



$$c_1 \frac{dT_1}{dt} = N_1 - H + F_{surf,1}$$


$$c_2 \frac{dT_2}{dt} = N_2 + H + F_{surf,2}$$

$$N_1 = \lambda_{1,1}T_1 + \lambda_{1,2}T_2 + F_{TOA,1} + F_{CO_2,1}$$

$$N_2 = \lambda_{2,1}T_1 + \lambda_{2,2}T_2 + F_{TOA,2} + F_{CO_2,2}$$

Rose and Rayborn 2016
Andrews and Webb 2017
Ceppi and Gregory 2017


Klein et al. 2017 Zhou et al. 2017 Dong et al. 2019



$$c_1 \frac{dT_1}{dt} = N_1 - H + F_{surf,1}$$

$$c_2 \frac{dT_2}{dt} = N_2 + H + F_{surf,2}$$


$$\overrightarrow{N} = \Lambda \overrightarrow{T} + \overrightarrow{F}_{TOA} + \overrightarrow{F}_{CO_2}$$



$$c_1 \frac{dT_1}{dt} = N_1 - H + F_{surf,1}$$

$$c_2 \frac{dT_2}{dt} = N_2 + H + F_{surf,2}$$

$$\overrightarrow{N} = \Lambda \overrightarrow{T} + \overrightarrow{F}_{TOA} + \overrightarrow{F}_{CO_2}$$




$$c_1 \frac{dT_1}{dt} = N_1 - H + F_{surf,1}$$


$$c_2 \frac{dT_2}{dt} = N_2 + H + F_{surf,2}$$


$$\overrightarrow{N} = \Lambda \overrightarrow{T} + noise$$


$$\overrightarrow{N} = \Lambda \overrightarrow{T} + noise$$




$$\overrightarrow{N} = \Lambda \overrightarrow{T} + noise$$












$$\overrightarrow{N} = \Lambda \overrightarrow{T} + noise$$



$$\overrightarrow{N} = \Lambda \overrightarrow{T} + noise$$

LongRunMIP, Rugenstein et al., 2019






$$\overrightarrow{N} = \Lambda \overrightarrow{T} + noise$$


 $Wm^{-2}K^{-1}$


Dong et al., 2019

 $Wm^{-2}K^{-1}$











$$\overrightarrow{N} = \Lambda \overrightarrow{T} + noise$$


Dong et al., 2019



$$\overrightarrow{N} = \Lambda \overrightarrow{T} + noise$$




$$\overrightarrow{N} = \Lambda \overrightarrow{T} + noise$$





$$\overrightarrow{N} = \Lambda \overrightarrow{T} + noise$$


evolution of abrupt4x feedback





evolution of abrupt4x feedback





































$$\overrightarrow{c} \cdot \frac{d\overrightarrow{T}}{dt} = \Lambda \overrightarrow{T} + \overrightarrow{F}_{TOA} - H + \overrightarrow{F}_{surf}$$


$$\overrightarrow{N}$$




Proistosescu et al 2018




Proistosescu et al 2018



Proistosescu et al 2018




Proistosescu et al 2018





$$\overrightarrow{c} \cdot \frac{d\overrightarrow{T}}{dt} = \Lambda \overrightarrow{T} + \overrightarrow{F}_{TOA} - H + \overrightarrow{F}_{surf}$$


$$\overrightarrow{N}$$





$$\overrightarrow{c} \cdot \frac{d\overrightarrow{T}}{dt} = \Lambda \overrightarrow{T} + \overrightarrow{F}_{TOA} - H + \overrightarrow{F}_{surf}$$














Conclusions

- Multiple regression method can be used to estimate spatial feedbacks from internal variability
- The MR method overestimates climate feedback due to anomalously high changes in TOA flux south of 30°S
- Overestimate could be due to "regression dilution" or nonlinearity in Green's functions - cause TBD